

# PERIODIC BILLIARD TRAJECTORIES IN SMOOTH CONVEX BODIES

R.N. KARASEV

**ABSTRACT.** We consider billiard trajectories in a smooth convex body in  $\mathbb{R}^d$  and estimate the number of distinct periodic trajectories that make exactly  $p$  reflections per period at the boundary of the body. In the case of prime  $p$  we obtain the lower bound  $(d-2)(p-1)+2$ , which is much better than the previous estimates.

## 1. INTRODUCTION

First, we give several definitions on billiards in convex bodies.

**Definition.** A *billiard trajectory* in a smooth convex body  $T \in \mathbb{R}^d$  is a polygon  $P \subset T$ , with all its vertices on the boundary of  $T$ , and at each vertex the direction of line changes according to the elastic reflection rule.

**Definition.** Let  $P$  be a periodic billiard trajectory. Its number of vertices is called its *length*.

From here on  $d$  will denote the dimension of convex body under consideration. We encode periodic trajectories by the sequence of their vertices  $(x_1, x_2, \dots, x_l)$ . The indices are considered mod  $l$ . We do not allow coincidences  $x_i = x_{i+1}$ , but  $x_i$  may equal  $x_j$  if  $i - j \neq 0, \pm 1$  mod  $l$ . Of course, the polygon segments can have mutual intersections.

The problem of finding lower bounds on the number of distinct trajectories of a given length has quite a long history. We mean by “distinct trajectories” the orbits of the dihedral group, acting naturally on trajectories.

The first lower bound was obtained in [1] for the two-dimensional case.

---

2000 *Mathematics Subject Classification.* 55M30.

*Key words and phrases.* billiard trajectories, Lyusternik-Snirelmann theory.

This research was supported by the Russian Foundation for Basic Research grants No. 03-01-00801 and 06-01-00648, and by the President of the Russian Federation grants No. MK-5724.2006.1 and No. MK-1005.2008.1.

In the case of length 2 and dimension  $d$  the paper [2] gives the lower bound  $d$ . This bound is tight, it may be seen by considering a generic ellipsoid.

The case of dimension 3 was considered in [5], but later an error was found, as noted in [8, 9].

In the book [7] the following problem was formulated (Problem 1.7):

**Problem.** *Let  $T \subset \mathbb{R}^d$  be a smooth convex body. Prove that it has at least  $d$  periodic billiard trajectories of length 3.*

In the papers [8, 9] some lower bounds were obtained for length  $n$ . There were two cases: odd  $n$  without other assumptions, and odd prime  $n$ . In the latter case the bounds were better. Here we give another lower bound for prime lengths.

**Theorem 1.** *Let  $T \subset \mathbb{R}^d$  be a smooth convex body,  $d \geq 3$ , let  $p > 2$  be a prime. Then there are at least  $(d - 2)(p - 1) + 2$  distinct periodic billiard trajectories of length  $p$  in  $T$ .*

The proof of this theorem is mainly based on results from [8, 9], calculating the cohomology ring of relevant configuration spaces. The case of non-prime length is not considered here.

## 2. COHOMOLOGICAL INDEX OF $Z_p$ -SPACES

In the sequel, if we want to consider  $Z_p$  as a group of transformations, we denote it  $G$ . If it is used as a ring of coefficients for cohomology it is denoted  $Z_p$ .

Consider the cohomology algebra  $A_G = H_G^*(\text{pt}, Z_p) = H^*(BG, Z_p)$ . It is well known [3] that  $A_G$  has two generators  $v, y$  of dimensions 1 and 2 respectively, the relations being

$$v^2 = 0, \quad \beta(v) = y.$$

We denote  $\beta(x)$  the Bockstein homomorphism. We see that  $A_G$  is  $Z_p$  in every dimension.

**Definition.** Let  $X$  be a  $G$ -space. The maximal  $n$  such that the natural map

$$H_G^n(\text{pt}, Z_p) \rightarrow H_G^n(X, Z_p)$$

is nontrivial, is called the *cohomological index* of  $X$  and denoted  $\text{hind } X$ .

Note that if the map  $H_G^n(\text{pt}, Z_p) \rightarrow H_G^n(X, Z_p)$  is trivial for some  $n$ , it should be trivial for all larger  $n$ , since every element of  $A_G$  can be obtained from any nonzero element of less dimension by multiplications by  $v, y$ , elements of  $Z_p^*$ , and by applying the Bockstein homomorphism.

Every  $G$ -space  $X$  gives a fibration

$$(X \times EG)/G \rightarrow BG.$$

If  $G$  acts on cohomology of  $X$  trivially, there exists a spectral sequence with  $E_2$ -term equal to  $H^*(X, Z_p) \otimes A_G$ , converging to  $H_G^*(X, Z_p)$ . If  $G$  has a nontrivial action on cohomology,  $E_2$ -term is  $E_2^{x,y} = H^x(G, H^y(X, Z_p))$ , where  $H^*(G, M) = \text{Ext}_{Z_p[G]}^*(M, Z_p)$  is the cohomology of a  $G$ -module  $M$ .

If  $G$  acts freely on  $X$ , we have  $H_G^*(X, Z_p) = H^*(X/G, Z_p)$ . In this case  $\text{hind } X$  does not exceed the dimension of  $X$ .

The following property is obvious by the definition of index:

**Lemma 1** (Monotonicity of index). *If there is a  $G$ -equivariant map  $X \rightarrow Y$ , then*

$$\text{hind } X \leq \text{hind } Y.$$

### 3. LYUSTERNIK-SCHNIRELMANN THEORY

We recall the definition of the Lyusternik-Schnirelmann category of a topological space.

**Definition.** *The relative category of a pair  $Y \subseteq X$  is the minimal cardinality of a family of contractible in  $X$  open subsets of  $X$ , covering  $Y$ . It is denoted  $\text{cat}_X Y$ . The number  $\text{cat}_X X = \text{cat } X$  is called the category of  $X$ .*

Here we state the Lyusternik-Schnirelmann theorem on the number of critical points in the form it is used in [8] for manifolds with boundary.

**Theorem 2.** *Let  $X$  be a compact smooth manifold with boundary. Let  $f : X \rightarrow \mathbb{R}$  be a smooth function, its gradient at  $\partial X$  always having strictly outward (inward) direction. Then  $f$  has at least  $\text{cat } X$  critical points.*

We will need the following lemma, that is a special case of the results of [6].

**Lemma 2.** *Let a space  $X$  have  $k$  cohomology classes  $x_1, \dots, x_k \in H^1(X, Z_p)$  and  $l$  classes of arbitrary positive dimension  $y_1, \dots, y_l \in H^*(X, Z_p)$ . If*

$$\beta(x_1)\beta(x_2)\dots\beta(x_k)y_1\dots y_l \neq 0,$$

*then  $\text{cat } X \geq 2k + l + 1$ .*

Actually, we use the following corollary of the previous lemma.

**Lemma 3.** *Let a finite group  $G$  act freely on a space  $X$ , assume that for some subgroup of prime order  $G' \subseteq G$  we have  $\text{hind}_{G'} X \geq n$ . Then  $\text{cat } X/G \geq n + 1$ .*

*Proof.* As it was noted in [9]  $\text{cat } X/G \geq \text{cat } X/G'$ .

$G'$  acts freely on  $X$ , hence  $H^*(X/G', Z_p) = H_{G'}^*(X, Z_p)$ . The latter algebra has a nonzero product of  $A_{G'}$  elements  $y^{\frac{n}{2}}$  or  $vy^{\frac{n-1}{2}}$ , for even or odd  $n$  respectively. So we can apply Lemma 2.  $\square$

#### 4. THE CONFIGURATION SPACE

Following the papers [8, 9] we describe the configuration spaces that arise naturally in the billiard problems. For a topological space  $X$  denote

$$G(X, p) = \{(x_1, \dots, x_p) \in X^p : x_1 \neq x_2, x_2 \neq x_3, \dots, x_{p-1} \neq x_p, x_p \neq x_1\}.$$

The space  $G(X, p)$  has an action of the dihedral group  $D_p$ , which has a subgroup of even permutations, isomorphic to  $Z_p = G$ .

Proposition 4.1 of [8] claims that  $G(\partial T, p)$  contains a  $D_p$ -invariant compact manifold with boundary  $G_\varepsilon(\partial T, p)$ , which is  $D_p$ -equivariantly homotopy equivalent to  $G(\partial T, p)$ .

In this section we find the cohomological  $G$ -index of  $G(\partial T, p)$ , obviously equal to the index of  $G_\varepsilon(\partial T, p)$ . The space  $\partial T$  is homeomorphic to a  $d - 1$ -sphere, so we write  $G(S^{d-1}, p)$ .

The first lower bound on the index of  $G(S^{d-1}, p)$  will be obtained by considering another space:  $G(\mathbb{R}^d, p)$ . Recall a special case of Proposition 2.2 from [8].

**Theorem 3.** *Let  $d \geq 2$ . The cohomology algebra  $H^*(G(\mathbb{R}^d, p), Z_p)$  is multiplicatively generated by  $d - 1$ -dimensional classes  $s_1, \dots, s_p$  and relations*

$$s_1^2 = s_2^2 = \dots = s_p^2 = 0, \quad s_1 s_2 \dots s_{p-1} + s_2 s_3 \dots s_p + s_3 s_4 \dots s_p s_1 + s_p s_1 \dots s_{p-2} = 0.$$

*The group  $G$  acts on  $s_1, \dots, s_p$  by cyclic permutations.*

Considering the action of  $G$  on  $G(\mathbb{R}^d, p)$ , we deduce a corollary.

**Theorem 4.** *If  $d \geq 2$ , then  $\text{hind } G(\mathbb{R}^d, p) = (d - 1)(p - 1)$ .*

*Proof.* Denote  $Y = G(\mathbb{R}^d, p)$ .

Consider a spectral sequence with  $E_2^{x,y} = H^x(G, H^y(Y, Z_p))$ . Note that  $H^*(Y, Z_p)$  are free  $Z_p[G]$ -modules in dimensions between 0 and  $(d - 1)(p - 1)$ . In dimension 0 it is  $Z_p$ , in dimension  $(d - 1)(p - 1)$  it is  $Z_p[G]/Z_p$ .

Let  $I_G^k$  be the ideal of  $A_G$ , consisting of elements of dimension at least  $k$ .

In  $E_2$ -term the bottom row is  $A_G$ , the top row is  $I_G^1$  with shifted by  $-1$  grading. The latter claim is deduced from the cohomology exact sequence for  $0 \rightarrow Z_p \rightarrow Z_p[G] \rightarrow Z_p[G]/Z_p \rightarrow 0$ .

All other rows of  $E_2$  have nontrivial groups in 0-th column only. Every differential of the spectral sequence is a homomorphism of  $A_G$ -modules. Hence, the intermediate rows cannot be mapped non-trivially to the bottom row. The only nonzero differential can map the top row to a part of the bottom row, isomorphic to  $I_G^{(d-1)(p-1)+1}$ . This map has to be nontrivial because the index of  $Y$  is finite.  $\square$

We see that if  $X$  contains  $\mathbb{R}^d$ , then  $G(X, p) \supseteq G(\mathbb{R}^d, p)$ . By the monotonicity of index we have  $\text{hind } G(X, p) \geq (d-1)(p-1)$ , in particular  $G(S^{d-1}, p) \geq (d-2)(p-1)$ .

Then we need more precise estimates on  $\text{hind } G(S^{d-1}, p)$ . Recall two theorems from [9] (Theorem 18, Theorem 19), describing the algebra  $H^*(G(S^{d-1}, p), Z_p)$ . We use the notation  $[n] = \{1, 2, \dots, n\}$ .

**Theorem 5.** *Let  $d \geq 4$  be even. Then  $H^*(G(S^{d-1}, p), Z_p)$  is generated by the elements*

$$u \in H^{d-1}(G(S^{d-1}, p), Z_p), \quad s_i \in H^{i(d-2)}(G(S^{d-1}, p), Z_p), i \in [p-2]$$

*and relations*

$$u^2 = 0, \quad s_i s_j = \frac{(i+j)!}{i!j!} s_{i+j}, \text{ if } i+j \leq p-2, \text{ or otherwise } s_i s_j = 0.$$

**Theorem 6.** *Let  $d \geq 3$  be odd. Then  $H^*(G(S^{d-1}, p), Z_p)$  is generated by the elements*

$$w \in H^{2d-3}(G(S^{d-1}, p), Z_p), \quad t_i \in H^{i(2d-4)}(G(S^{d-1}, p), Z_p), i \in \left[ \frac{p-3}{2} \right]$$

*and relations*

$$w^2 = 0, \quad t_i t_j = \frac{(i+j)!}{i!j!} t_{i+j}, \text{ if } i+j \leq \frac{p-3}{2}, \text{ or otherwise } t_i t_j = 0.$$

Theorem 5 is formulated in [9] for the cohomology coefficients  $\mathbb{Q}$ . But the proof is valid without change for  $Z_p$ , since we only have to divide by  $i!$ , where  $i \leq p-2$ .

Note that  $G$  acts on the cohomology in these theorems trivially, since the cohomology is either  $Z_p$  or 0 in every dimension.

Finally, let us give another lower bound for  $\text{hind } G(S^{d-1}, p)$ . We have a  $D_p$ -equivariant map from the Stiefel variety of 2-frames in  $\mathbb{R}^d$   $V_d^2 \rightarrow G(S^{d-1}, p)$ , defined as follows. Take some regular  $p$ -gon in the plane. Every frame  $(e_1, e_2) \in V_d^2$  gives an embedding of this  $p$ -gon to

$S^{d-1}$ , which is  $D_p$ -equivariant. So  $\text{hind } G(S^{d-1}, p) \geq \text{hind } V_d^2$ , the latter index being equal to (see [4])  $2d - 3$ .

**Theorem 7.** *If  $d \geq 3$ , then  $\text{hind } G(S^{d-1}, p) = (d - 2)(p - 1) + 1$ .*

*Proof.* First, as it was mentioned in [5, 8], the Morse theory shows that  $G(S^{d-1}, p)$  is homotopy equivalent to a CW-complex of dimension  $(d - 2)(p - 1) + 1$ . So the index is naturally bounded from above and we prove the lower bound only.

If  $p = 3$   $(p - 1)(d - 2) + 1 = 2d - 3$  and the lower bound is already done. So we consider the case  $p \geq 5$ .

Note that  $H^*(G(S^{d-1}, p), Z_p)$  is multiplicatively generated by  $(u, s_1)$  and  $(w, t_1)$  for even and odd  $d$  respectively.

First consider the case of even  $d$ .

Consider the spectral sequence with  $E_2 = H^*(G(S^{d-1}, p), Z_p) \otimes A_G$ . In this case the multiplicative generators of  $H^*(G(S^{d-1}, p), Z_p)$  have dimension at most  $d - 1$ . If the differentials  $d_m$  were trivial for  $m \leq d$ , then they should be trivial for  $m > d$  from the dimension of generators. But in this case  $\text{hind } G(S^{d-1}, p) = +\infty$ , which is false. So some of  $d_m$  is nontrivial for  $m \leq d$ .

Note that in  $E_2$  the bottom row is  $A_G$ , the next from bottom is  $s_1 A_G$ , then  $u A_G$ . Let  $d_m$  be the first nontrivial differential. The generators  $u$  and  $s_1$  cannot be mapped non-trivially to the bottom row, since in this case  $\text{hind } G(S^{d-1}, p)$  would be at most  $d - 1$ . The only possibility for nontrivial  $d_m$  is  $d_2$  that has  $d_2(u) = a s_1 y$  ( $a \in Z_p^*$ ). In this case  $E_3$  will be multiplicatively generated by  $v, y \in A_G$ , the image of  $s_1$  and of  $u s^{p-2}$ , denote the latter by  $z$ . The relations would be (besides those of  $A_G$ ):

$$s_1^{p-1} = 0, \quad s_1 y = 0, \quad z^2 = 0.$$

That means that  $E_3$  has two full rows in the top and in the bottom, and several rows in between contain  $s_1^k$  and  $s_1^k v$  for  $k \in [p - 2]$ .

The next  $d_m$  cannot map  $z$  non-trivially if  $m \leq (p - 1)(d - 2) + 1 = \dim z$ . They also cannot map  $s_1$  non-trivially from the dimension considerations and the lower bound for index. The latter statement can also be deduced from the fact that all differentials are homomorphisms of  $A_G$ -modules. So finally  $d_{(p-1)(d-2)+2}$  has to map  $z$  to the bottom row and we have  $\text{hind } G(S^{d-1}, p) = (p - 1)(d - 2) + 1$ .

Now consider the case of odd  $d$ . Everything is quite the same, but besides the alternative that gives  $\text{hind } G(S^{d-1}, p) = (p - 1)(d - 2) + 1$  we have another alternative. It could happen that  $d_2$  is trivial and the first nontrivial  $d_{2d-2}$  maps  $w$  non-trivially to the bottom row. But in this case  $\text{hind } G(S^{d-1}, p) = 2d - 3$ , which is less then the bound  $(d - 2)(p - 1)$  for  $p \geq 5$ .  $\square$

## 5. PROOF OF THEOREM 1

Consider the function on  $G(\partial T, p)$

$$f : (x_1, \dots, x_p) \mapsto \sum_{i=1}^p |x_i x_{i+1}|.$$

If  $\frac{\partial f}{\partial x_i} = 0$ , then at  $x_i$  the polyline  $x_1 x_2 \dots x_p x_1$  reflects by the elastic reflection rule.  $f$  can be considered as a function on  $G(\partial T, p)/D_p$ , its critical points being in one-to-one correspondence with distinct periodic billiard trajectories of length  $p$ .

If  $T$  is not strictly convex, some line segments of the trajectory can lie on  $\partial T$ . But in this case all the segments must lie on the same line, which is impossible.

Following [8] we consider  $G_\varepsilon(\partial T, p)$  instead of  $G(\partial T, p)$  and apply Theorem 2 to  $f$ .

Then we estimate  $\text{cat } G(\partial T, p)/D_p \geq (d-2)(p-1)+2$  by Theorem 7 and Lemma 3.

## 6. CONCLUSION

For an arbitrary  $d$ -manifold  $M$  ( $d \geq 2$ )  $\text{hind } G(M, p)$  is between  $(d-1)(p-1)$  (Theorem 4) and  $(d-1)(p-1)+1$  (by the dimension considerations). It seems probable that for closed  $M$   $\text{hind } G(M, p) = (d-1)(p-1)+1$ , as it is for spheres (Theorem 7).

The author would like to thank A. Yu. Volovikov for useful discussions on the subject of the paper.

## REFERENCES

- [1] G. Birkhoff. Dynamical systems. // Amer. Math. Soc. Coll. Publ., 9, 1927
- [2] N.H. Kuiper. Double normals of convex bodies. // Israel Journal of Mathematics, 2, 1964, 71–80
- [3] Wu Yi Hsiang. Cohomology theory of topological transformation groups. Berlin-Heidelberg-New-York: Springer Verlag, 1975
- [4] V.V. Makeev. Knaster's problem and almost spherical sections. // Math. Sbornik, 180(3), 1989, 424–431
- [5] I.K. Babenko. Periodic trajectories in three-dimensional Birkhoff billiards. // Math. Sbornik, 181(9), 1990, 1155–1169
- [6] E. Fadell, S. Husseini. Category weight and Steenrod operations. // Boletín de la Sociedad Matemática Mexicana, 37, 1992, 151–161
- [7] V. Klee, S. Wagon. Old and new unsolved problems in plane geometry and number theory. Dolciani Mathematical Expositions, The Mathematical Association of America, 1996

- [8] M. Farber, S. Tabachnikov. Topology of cyclic configuration spaces and periodic trajectories of multi-dimensional billiards. // Topology, 41(3), 2002, 553–589
- [9] M. Farber. Topology of billiard problems, II. // Duke Mathematica Journal, 115, 2002, 587–621

*E-mail address:* `r_n_karasev@mail.ru`

ROMAN KARASEV, DEPT. OF MATHEMATICS, MOSCOW INSTITUTE OF PHYSICS  
AND TECHNOLOGY, INSTITUTSKIY PER. 9, DOLGOPRUDNY, RUSSIA 141700